



Contents lists available at ScienceDirect

## Contact Lens and Anterior Eye

journal homepage: [www.elsevier.com/locate/clae](http://www.elsevier.com/locate/clae)

## Contact Lens Evidence-Based Academic Reports (CLEAR)

James S. Wolffsohn<sup>a,\*</sup>, Philip B. Morgan<sup>b</sup>, Melissa Barnett<sup>c</sup>, Laura E. Downie<sup>d</sup>,  
Deborah S. Jacobs<sup>e,f</sup>, Lyndon Jones<sup>g,h</sup>, Kathryn Richdale<sup>i</sup>, Fiona Stapleton<sup>j</sup>,  
Stephen J. Vincent<sup>k</sup>, Mark Willcox<sup>j</sup>

<sup>a</sup> Optometry and Vision Science, Aston University, Birmingham, UK

<sup>b</sup> Eurolens Research, Division of Pharmacy and Optometry, University of Manchester, UK

<sup>c</sup> University of California, Davis Eye Center, Sacramento, CA, USA

<sup>d</sup> Department of Optometry and Vision Sciences, The University of Melbourne, Australia

<sup>e</sup> Massachusetts Eye & Ear, Cornea and Refractive Surgery Service, Boston, MA, USA

<sup>f</sup> Harvard Medical School, Boston, MA, USA

<sup>g</sup> Centre for Ocular Research & Education (CORE), School of Optometry & Vision Science, University of Waterloo, Waterloo, Canada

<sup>h</sup> Centre for Eye and Vision Research (CEVR), Hong Kong

<sup>i</sup> College of Optometry, University of Houston, Houston, TX, USA

<sup>j</sup> School of Optometry and Vision Science, University of New South Wales, Sydney, NSW, Australia

<sup>k</sup> Queensland University of Technology (QUT), Centre for Vision and Eye Research, School of Optometry and Vision Science, Contact Lens and Visual Optics Laboratory, Brisbane, Australia

## ARTICLE INFO

## Keywords:

Abbreviations

Contact lens evidence-based academic reports

(CLEAR)

Overnight wear

Rigid corneal lens

Since contact lenses were invented in 1887, innovations have included advances in optical design, material, care systems, wear modality, lens size, lens shape and applications. Over 19,000 peer reviewed academic papers on the contact lenses have been published. The Contact Lens Evidence-based Academic Reports (CLEAR) follow the exemplary work of organisations such as the Tear Film and Ocular Surface Society dry eye [1], meibomian gland disease [2], contact lens discomfort [3] and dry eye II [4] workshops and the International Myopia Institute white papers on myopia control [5], in collating and appraising the academic literature in an ocular field of interest. CLEAR represent the work of nearly 100 multidisciplinary experts in the field, who set out to critically review, synthesise and summarise the research evidence on contact lenses to date; this serves to inform both clinical practice, manufacturing innovation and future research directions.

## 1. The process

CLEAR was conceived by James Wolffsohn in June 2019 and the British Contact Lens Association (BCLA) executive committee gave their backing in September 2019. Philip Morgan was elected as executive vice-chair and the name for the initiative, report topics and leading experts as chairs (Table 1) were agreed together with Cheryl Donnelly, the then Chief Executive Officer of the BCLA. Sponsors were sought to cover the costs of production and publication, but had no input concerning the scope or content of the reports.

Interested clinicians and scientists were invited to apply to working groups and experts in the field (identified by the BCLA and report chairs) were selected to contribute to one of the reports that best fitted their area of expertise and/or practice. An inclusive approach was adopted, while limiting the number of participants from any one research group or company on any single report to ensure a balanced representation.

The report committees, led by their chair, developed an outline of the

\* Corresponding author.

E-mail address: [j.s.w.wolffsohn@aston.ac.uk](mailto:j.s.w.wolffsohn@aston.ac.uk) (J.S. Wolffsohn).

<https://doi.org/10.1016/j.clae.2021.02.011>

Available online 25 March 2021

1367-0484/© 2021 British Contact Lens Association. Published by Elsevier Ltd. All rights reserved.

**Table 1**  
CLEAR Reports, their Chairs and their Harmonisers.

CLEAR Report	Chair	Harmoniser
Anatomy and physiology of the anterior eye [6]	Laura Downie	James Wolffsohn
Contact lenses wettability, cleaning, disinfection and interactions with tears [7]	Mark Willcox	Lyndon Jones
Effect of contact lens materials and designs on the anatomy and physiology of the eye [8]	Philip Morgan	Mark Willcox
Contact lens optics [9]	Kathryn Richdale	Philip Morgan
Orthokeratology [10]	Stephen Vincent	Lyndon Jones
Scleral lenses [11]	Melissa Barnett	Philip Morgan
Medical use of contact lenses [12]	Deborah Jacobs	Fiona Stapleton
Contact lens complications [13]	Fiona Stapleton	James Wolffsohn
Evidence-based contact lens practice [14]	James Wolffsohn	Fiona Stapleton
Contact lens technologies of the future [15]	Lyndon Jones	Mark Willcox

subtopics to be covered in their report in January and February 2020 and these were reviewed in March to minimise gaps and overlap. The chairs allocated the subsections of their report to members of their committee and writing commenced in April; drafts were returned to chairs by the end of May to collate and review. This version was reviewed and edited by all committee members by August 2020. From August to September the refined reports were sent to all CLEAR members for review. Their comments were sent back to chairs for addressing (in a similar fashion, but more extensively than would occur through a typical peer review academic journal process) by November 2020. Simultaneously, a medical illustrator was commissioned to draw the necessary original artwork for the reports. The harmonisers (Table 1) met (virtually due to COVID-19 travel limitations) to agree on the standardisation of terminology, abbreviations, formatting, remaining areas of report overlap and abstract development. The finalised reports were submitted to the BCLA's journal, *Contact Lens and Anterior Eye*, in January 2021.

Evidence-based practice can be defined as the “conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients” [16]. It involves integrating the best available, clinically relevant, scientific research evidence with a clinician's expertise and an individual patient's values and environment (see CLEAR Evidence-based Practice Report) [14]. Evaluating scientific research findings and using them to make the best clinical decision for patients is a key aim of all evidence-based practice, including fitting and managing contact lenses. The quality of research evidence generally comes from the study design [14]. Formal risk of bias tools also exist to assist clinicians with appraising the quality of an individual study rather than simply relying on the evidence level [17].

Larger cohort studies are considered more informative than expert opinion, case reports or case series and are generally used for epidemiological studies. For management decisions, randomised controlled trials are considered to provide the highest level of evidence-basis; they limit unconscious bias through masking (ideally of the participants and clinical researchers) as to which treatment participants are provided, and randomise participants to treatment and control (or placebo) options. Systematic reviews collate, appraise and synthesise evidence from multiple papers that fit pre-specified eligibility criteria, to answer a specific research question. Hence they are generally considered a higher

level of evidence than individual research papers and are often used to inform professional clinical guidelines, which interpret and contextualise their finding to guide and regulate clinical practice. A number of international, evidence-based reviews that inform elements of contact lens practice have been conducted recently, based on a consensus-building workshop approach [5,18–20].

Individual studies within a given level of the hierarchy level (such as randomised controlled trials) may differ in their ‘quality’, due to differences in their study design, tests performed, cohort selection and participant numbers. Some contact lens research employs study designs that are not explicitly described in hierarchical models of research quality or common in general medicine; these include cross-over, contralateral and monadic designs which are often used to understand the clinical performance of different lenses and care products [14]. While systems for rating the level of evidence of individual papers or hypotheses have been developed, they are not robust enough to systematically apply [21], hence the approach taken in CLEAR is to present summaries of the research findings and to critically appraise the evidence on relevant topics.

## 2. Approach to terminology

Inconsistent terminologies have developed within the field of contact lenses that can be confusing to students, clinicians, researchers and other stakeholders. Several anatomical terms, named after individuals, such as Bowman's and Descemet's membranes, have been renamed the anterior and posterior limiting lamina by the Federative Committee on Anatomical Terminology (FCAT) [22] and these have been adopted by CLEAR. Whilst the FCAT also renamed ‘Meibomian glands’ as ‘tarsal glands’, this terminology does not logically follow other terms in the field. Meibomian glands produce meibum, and the term tarsal is only attributed to a single anatomical ‘plate’ that is not the sole location of the glands; lacrimal glands are named after the secretion they produce and not their location. Hence, ‘meibomian’ gland terminology, also used in the dry eye disease literature, has been retained in the CLEAR reports. Likewise, tarsal conjunctiva is a region of the palpebral conjunctiva rather than an appropriate term to describe this tissue on the underside of the eyelids.

The term rigid gas permeable lenses (RGP) was developed to differentiate the first oxygen-permeable hard/rigid lenses from earlier oxygen impermeable materials, such as poly methyl methacrylate (PMMA). In more recent times, this has been truncated to simply ‘gas permeable’ or ‘GP’ by some authors as ‘rigid’ was felt to suggest to potential patients that these lenses would cause discomfort [23]. However, all modern contact lenses (soft or rigid) are ‘gas permeable’ and scleral lenses are also RGPs, yet the term is generally used to describe exclusively corneal lenses. Hence, a poll of CLEAR members was conducted and 62 % were in favour of adopting a change in terminology to ‘corneal lens’, 18 % against (mainly as they felt soft lenses also ‘landed’ on the cornea) and the rest (21 %) were equivocal. Since their ‘rigidity’ is a key feature of the optical and health benefits of these lenses, the term ‘rigid corneal lens’ was adopted throughout CLEAR. Scleral lens terminology has recently been redefined [24] and CLEAR has accepted this approach, and thus all rigid lenses that vault the cornea are termed ‘scleral lenses’. While regulatory terminology denoted extended wear as 7 days and 6 nights, and continuous wear as up to 30 days and 29 nights [25], these definitions overlap and are used interchangeably in the literature. Research suggests that there are no marked clinical differences between these modalities [26]. Hence the terminology ‘planned’ or ‘sporadic’ ‘overnight wear’ is more appropriate for clinical use and has been adopted in these reports.

The CLEAR harmonisers carefully considered the use of abbreviations throughout the reports, using the principles articulated in Cochrane reviews [27] that they should be used sparingly and only if they are widely known across the broad readership, are used frequently and enhance readability. Two word abbreviations were only adopted where the abbreviation is used more commonly than the words they represent. It is hoped this general list of terms will assist standardisation in future publications in the field and to support new eye care practitioners. Standard unit and country abbreviations are not articulated in full due to these being commonly accepted terms.

General list of abbreviations for the field of contact lenses->

BAK	benzalkonium chloride
BOZR/BOZD	back optic zone radius/diameter
CIE	Corneal infiltrative event
CLD	contact lens discomfort
CLIDE	contact lens induced dry eye
CLPC	contact lens-induced papillary conjunctivitis
Dk/t	oxygen permeability/transmissibility
ECP	eye care practitioner
EDOF	extended depth of focus
EDTA	ethylenediaminetetraacetic Acid
HEMA	2-hydroxyethyl methacrylate
HPMC	hydroxypropyl methylcellulose
HVID	horizontal visible iris diameter
LIPCOF	lid-parallel conjunctival folds
LWE	lid wiper epitheliopathy
MGD	meibomian gland dysfunction
MK	microbial keratitis
MPDS	multipurpose disinfecting solution
Ortho-k	orthokeratology
PEG	polyethylene glycol
PHMB	polyhexamethylene biguanide
PMMA	polymethyl methacrylate
PolTF	post-lens tear film
PVA	polyvinyl alcohol
PVP	polyvinyl pyrrolidone
SICS	solution induced corneal staining
SiHy	silicone-hydrogel soft contact lens
VPA	vertical palpebral aperture

3. CLEAR

The collaboration between experts in the field of contact lenses and the anterior eye has been inspiring and productive, despite the enforced ‘virtual’ nature of the interactions due to the COVID-19 pandemic. These reports bring together the evidence, and consensus where this was lacking, to inform clinical practice, identifying areas where further research is needed and determining where there are opportunities for new innovations from industry.

Acknowledgements

The CLEAR initiative was facilitated by the BCLA, with financial support by way of Educational Grants for collaboration, publication and dissemination provided by Alcon and CooperVision.

References

- [1] Introduction to the report of the international Dry Eye WorkShop. *Ocular Sci* 2007; 5:69–70.
- [2] Nichols KK. The international workshop on meibomian gland dysfunction: introduction. *Invest Ophthalmol Vis Sci* 2011;52:1917–21.
- [3] Nichols JJ, Jones L, Nelson JD, Stapleton F, Sullivan DA, Willcox MD, et al. The TFOS international workshop on contact lens discomfort: introduction. *Invest Ophthalmol Vis Sci* 2013;54:TFOS1–6.
- [4] Nelson JD, Craig JP, Akpek EK, Azar DT, Belmonte C, Bron AJ, et al. TFOS DEWS II introduction. *Ocul Surf* 2017;15:269–75.
- [5] Wolffsohn JS, Flitcroft DI, Gifford KL, Jong M, Jones L, Klaver CCW, et al. IMI – myopia control reports overview and introduction. *Invest Ophthalmol Vis Sci* 2019; 60:M1–19.
- [6] Downie LE, Bandlitz S, Bergmanson JPG, Craig JP, Dutta D, Maldonado-Codina C, et al. CLEAR – anatomy and physiology of the anterior eye. *Contact Lens Anterior Eye* 2021;44(2):132–56.
- [7] Willcox M, Keir N, Maseedupally V, Masoudi S, McDermott A, Mobeen R, et al. CLEAR – contact lenses wettability, cleaning, disinfection and interactions with tears. *Contact Lens Anterior Eye* 2021;44(2):157–91.
- [8] Morgan P, Murphy PJ, Gifford K, Gifford P, Golebiowski B, Johnson L, et al. CLEAR – effect of contact lens materials and designs on the anatomy and physiology of the eye. *Contact Lens Anterior Eye* 2021;44(2):192–219.
- [9] Richdale K, Cox I, Kollbaum P, Bullimore MA, Bakaraju RC, Gifford P, et al. CLEAR – contact lens optics. *Contact Lens Anterior Eye* 2021;44(2):220–39.
- [10] Vincent SJ, Cho P, Chan KY, Fadel D, Ghorbani-Mojarrad N, González-Mejome JM, et al. CLEAR – orthokeratology. *Contact Lens Anterior Eye* 2021;44(2):240–69.
- [11] Barnett M, Courey C, Fadel D, Lee K, Michaud L, Montani G, et al. CLEAR scleral lenses. *Contact Lens Anterior Eye* 2021;44(2):270–88.
- [12] Jacobs DS, Carrasquillo KG, Cottrell PD, Fernández-Velázquez FJ, Gil-Cazorla R, Jalbert I, et al. CLEAR – medical use of contact lenses. *Contact Lens Anterior Eye* 2021;44(2):289–329.
- [13] Stapleton F, Bakkar M, Carnt N, Chalmers R, Kumar A, Marasini S, et al. CLEAR – contact lens complications. *Contact Lens Anterior Eye* 2021;44(2):330–67.
- [14] Wolffsohn JS, Dumbleton K, Huntjens B, Kandel H, Koh S, Kunnen CME, et al. CLEAR – evidence based contact lens practice. *Contact Lens Anterior Eye* 2021;44(2):368–97.
- [15] Jones L, Hui A, Phan C-M-M, Read ML, Azar D, Buch J, et al. CLEAR – contact lens technologies of the future. *Contact Lens Anterior Eye* 2021;44(2):398–430.
- [16] Sackett DL, Rosenberg WMC, Gray JAM, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *Br Med J* 1996;312:71–2.
- [17] Featherston R, Downie LE, Vogel AP, Galvin KL. Decision making biases in the allied health professions: a systematic scoping review. *PLoS One* 2020;15: e0240716.
- [18] Craig JP, Nelson JD, Azar DT, Belmonte C, Bron AJ, Chauhan SK, et al. TFOS DEWS II report executive summary. *Ocul Surf* 2017;15:802–12.
- [19] Nichols KK, Foulks GN, Bron AJ, Glasgow BJ, Dogru M, Tsubota K, et al. The international workshop on meibomian gland dysfunction: executive summary. *Invest Ophthalmol Vis Sci* 2011;52:1922–9.
- [20] Nichols JJ, Willcox MD, Bron AJ, Belmonte C, Ciolino JB, Craig JP, et al. The TFOS international workshop on contact lens discomfort: executive summary. *Invest Ophthalmol Vis Sci* 2013;54:7–13.
- [21] Atkins D, Eccles M, Flottorp S, Guyatt GH, Henry D, Hill S, et al. Systems for grading the quality of evidence and the strength of recommendations I: critical appraisal of existing approaches The GRADE Working Group. *BMC Health Serv Res* 2004;4:38.
- [22] Terminology. FICoA. Terminologia histologica: international terms for human cytology and histology. 2008.
- [23] Efron N. Obituary—rigid contact lenses. *Cont Lens Anterior Eye* 2010;33:245–52.
- [24] Michaud L, Lipson M, Kramer E, Walker M. The official guide to scleral lens terminology. *Cont Lens Anterior Eye* 2020;43:529–34.
- [25] Riley C, Chalmers RL. Survey of contact lens-wearing habits and attitudes toward methods of refractive correction: 2002 versus 2004. *Optom Vis Sci* 2005;82: 555–61.
- [26] Bialasiewicz AA. Infection immunology in silicone hydrogel contact lenses for continuous wear—a review. *Klin Monbl Augenheilkd* 2003;220:453–8.
- [27] Cochrane. Cochrane style manual. 2020.